

Cluster states of Fermions in the single l -shell model

András Csordás*

*Research Group for Statistical Physics of the Hungarian Academy of Sciences,
Pázmány P. Sétány 1/A, H-1117 Budapest, Hungary*

Éva Szőke†

*Department of Physics of Complex Systems, Eötvös University,
Pázmány P. Sétány 1/A, H-1117 Budapest, Hungary‡*

Péter Szépfalussy§

*Department of Physics of Complex Systems, Eötvös University,
Pázmány P. Sétány 1/A, H-1117 Budapest, Hungary† and
Research Institute for Solid State Physics and Optics, P. O. Box 49, H-1525 Budapest, Hungary
(Dated: February 2, 2008)*

The paper concerns the ground state structure of the partly filled l -shell of a fermionic gas of atoms of spin s in a spherically symmetric spin independent trap potential. At particle numbers $N = n(2s + 1)$, $n = 1, 2, \dots, 2l + 1$ the basic building blocks are clusters consisting of $(2s + 1)$ atoms, whose wave functions are completely symmetric and antisymmetric in space and spin variables, respectively. The creation operator of a cluster is constructed whose repeated application to the vacuum leads to the multi-cluster state. Ground state energy expressions are derived for the n -cluster states at different l, s values and interpreted in simple terms.

PACS numbers: 05.30.Fk, 31.15.Hz, 21.60.Cs, 74.20.Fg

I. INTRODUCTION

The many body problem as applied to finite systems has a long history in atomic and nuclear physics [1, 2, 3, 4]. One of the central problems has been the nature of the ground state in case of a partially filled shell.

When we consider atoms (a gas at zero temperature) in an external potential the possible behaviors are quite rich. One can assume that the collision between atoms does not excite internal degrees of freedom and the atoms can be regarded as structureless objects whose spins are fixed being in a definite hyperfine state. The external potential can be supplied by a magnetic or optical trap [5]. In the past few years a very intensive research has been continued in case of Bose-particles both theoretically and experimentally in a variety of such systems. Fermion systems along these lines have been less studied until recently, but important achievements have been already available and one can be sure that rapid development will continue in the future. In particular the achievement and study of the superfluid state has become one of the frontiers in physics (see for Reviews [6, 7, 8, 9], which contain references to earlier works).

In this paper we treat fermionic systems. The external potential will be assumed to be spherically symmet-

ric and the interaction between atoms will be described by a spin independent δ function like attractive pseudopotential. Within the single l -shell model of fermionic atoms with arbitrary large spins we study the ground state properties of the gas.

We concentrate on cluster states containing particles of numbers $N_n = n(2s + 1)$ where s is the spin of the particles and $n = 1, 2, \dots, 2l + 1$. The existence of such clusters have been pointed out by us previously [10]. In the present paper we extend the investigations in several directions. It is advantageous to use the second quantized representation, which makes possible, among other things, to obtain the multi-cluster states by repeated applications of the cluster creation operator to the vacuum. Besides numerical calculations extensive analytical studies are carried out. It is shown that to a good approximation the hamilton operator can be replaced by a model one built of the the operator of the particle number, of the quadratic Casimir operators of the groups $SU(2l + 1)$ and $SO(2l + 1)$. It is shown that within the model the clusters can be conceived as interacting (Cooper) pairs if $s > 1/2$, the interaction being of statistical origin. The model Hamiltonian coincides with the true one for $l = 1, 2$ (The special cases mainly investigated in [10]).

For the model hamiltonian the cluster states are explicitly given and it is pointed out that the energy of the n -cluster state can be written as the sum of energies of the clusters proportional to n and an “interaction term” term between the clusters proportional to $n(n - 1)/2$. It is found that the second term disappears when $s = 1/2$. For $s = 1/2$ it was shown by Racah in 1952 [11] that the ground state consists of independent pairs.

*Electronic address: csordas@tristan.elte.hu

†Electronic address: jeva@ludens.elte.hu

‡The work has been done while beeing a member of the teaching staff of the Budapest University Catholic Secondary School, Szabó Ilonka u. 2-4, H-1015 Budapest, Hungary

§Electronic address: psz@complex.elte.hu

Investigations of particles in an open shell has been an important area in atom and nuclear physics. As discussed above in the trapped gas of Fermi atoms new features appear due to the fact that the spin of the particles can be higher than $1/2$. Comparing with the situation in the atom shell a further important difference is that instead of the long ranged Coulomb force between the electrons the atom-atom interaction which has to be considered is short ranged, while comparing the situation with that of an open neutron shell in a nucleus an important difference is that in our case the open shell is an l -one (since no-spin orbit interaction is present) as contrasted to the j -one in the nucleus.

Though our main purpose in this paper is to enlarge the picture we have about the dynamics of fermionic particles in a partly filled shell, a few words about the relevance of the model for physically realizable situations in case of trapped gases are in order. Obviously optical traps are the suitable ones which allow the free rotation of the spins. It is assumed that the external potential contains besides the confining potential (which is typically of a harmonic oscillator type) the mean field of the atoms building the closed shells. There are two conditions then to be fulfilled in order that the single- l shell model apply. Firstly, the mean field due to the closed shells should be strong enough that the possible degeneracy of the levels of different l values be lifted considerable. Secondly, the characteristic interaction energy of the atoms in the partly occupied shell should be smaller than the relevant level distance in the external potential. Both requirements can be satisfied simultaneously if the strength of the interaction between the atoms is weak and the number of the atoms in the trapped gas is sufficiently large [12, 13].

The paper is organized as follows. In Section II the Hamiltonian is written in terms of irreducible tensor operators and a model Hamiltonian is introduced built from the operator of particle number and from the quadratic Casimir operators of the groups $SU(2l+1)$ and $SO(2l+1)$. The Section III is devoted to the extensive investigation of the one-cluster state. The symmetrizer playing an important role in creating the cluster states is analyzed in detail in Section IV. Energy eigenvalues of the model Hamiltonian for cluster states generated by the cluster creation operator are calculated in Section V. It is shown in Section VI that the expectation value of the Hamiltonian agrees with that of the model one for cluster states. Section VII contains the summary and a discussion of the results. Appendices A and B present certain steps of proofs outlined in the main text. Appendix C discusses the special case when the spin of the particles is $1/2$.

II. FORMULATION

Let us assume that the trap potential is spherically symmetric. At zero temperature the open shell consists

of fermions of spin s which partially fill the (n, l) shell. The one particle normalized wave functions are given by

$$\Psi_{n,l,m,s,\nu}(r, \vartheta, \varphi, \sigma) = R_{n,l}(r) Y_m^l(\vartheta, \varphi) \chi_\nu^s(\varsigma), \quad (1)$$

Here ς is the discrete spin variable and $\chi_\nu^s(\varsigma)$ is the normalized spin eigenfunction. In the model the quantum numbers (n, l, s) are fixed, m and ν can take the values $m = -l, -l+1, \dots, l$ and $\nu = -s, -s+1, \dots, s$ and the particle number N can vary between 0 and $(2l+1)(2s+1)$. The functions (1) are eigenfunctions of the one-particle Hamiltonian, which contains the kinetic energy term, the trap potential and the averaged field potential of the closed shells. The interaction between the particles in the partially filled shell is given by

$$H_{int} = -\frac{\lambda}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^N \delta(\mathbf{r}_i - \mathbf{r}_j), \quad \lambda > 0 \quad (2)$$

corresponding to a spin independent s -wave scattering with negative scattering length. Our aim is to diagonalize (2) on the fixed basis (1).

Let us denote the operator which annihilates a particle with quantum numbers (n, l, m, s, ν) by $a_{m,\nu}$. The Hamiltonian (2) in second quantization reads as

$$\begin{aligned} \hat{H}_{int} &\equiv \hat{h}E_0/\pi = -\frac{E_0}{2} \sum_{\substack{m_1, m_2 \\ m_3, m_4}} \sum_{\nu_1, \nu_2} f_{m_1, m_2; m_3, m_4} \\ &\times a_{m_1, \nu_1}^+ a_{m_2, \nu_2}^+ a_{m_4, \nu_2} a_{m_3, \nu_1}, \end{aligned} \quad (3)$$

where E_0 is the characteristic energy

$$E_0 = \lambda \int_0^\infty |R_{n,l}(r)|^4 r^2 dr, \quad (4)$$

and f can be expressed in terms of the Wigner- $3j$ symbols [14] in two equivalent forms

$$\begin{aligned} f_{m_1, m_2; m_3, m_4} &= \int d\Omega Y_{m_1}^{l*}(\Omega) Y_{m_2}^{l*}(\Omega) Y_{m_3}^l(\Omega) Y_{m_4}^l(\Omega) \\ &= \frac{[l]^2}{4\pi} \sum_{L=0}^{2l} [L] \begin{pmatrix} L & l & l \\ 0 & 0 & 0 \end{pmatrix}^2 \\ &\times \sum_{M=-L}^L \begin{pmatrix} l & l & L \\ m_1 & m_2 & -M \end{pmatrix} \begin{pmatrix} l & l & L \\ m_3 & m_4 & -M \end{pmatrix} \quad (5) \\ &= \frac{[l]^2}{4\pi} \sum_{L=0}^{2l} [L] \begin{pmatrix} L & l & l \\ 0 & 0 & 0 \end{pmatrix}^2 (-1)^{(m_2+m_3)} \\ &\times \sum_{M=-L}^L \begin{pmatrix} l & l & L \\ m_1 & -m_3 & -M \end{pmatrix} \begin{pmatrix} l & l & L \\ m_4 & -m_2 & -M \end{pmatrix} \end{aligned}$$

For notational simplicity we introduced the symbol $[\dots]$ defined by

$$[p] \equiv (2p+1).$$

The dimensionless Hamiltonian $\hat{h}$ (see Eq. (3)) can be written as

$$\hat{h} = \frac{[l]}{8} \hat{N} - \frac{[l]^2}{8} \sum_{L=0}^{2l} \begin{pmatrix} l & l & L \\ 0 & 0 & 0 \end{pmatrix}^2 \hat{B}_L^2, \quad (7)$$

where

$$\hat{B}_L^2 = \sum_{M=-L}^L (-1)^{L-M} \hat{B}_{L,M} \hat{B}_{L,-M} \quad (8)$$

(in eq. (7) only the terms L even remain). The operators $\hat{B}_{L,M}$ defined as

$$\begin{aligned} \hat{B}_{L,M} &= \sum_{m=-l}^l \sum_{\nu=-s}^s (-1)^{l-m} \sqrt{[L]} \\ &\times \begin{pmatrix} l & l & L \\ m & M-m & -M \end{pmatrix} a_{m,\nu}^+ a_{m-M,\nu} \end{aligned} \quad (9)$$

are spin scalars and irreducible tensoroperators with respect to angular momentum [1]. Special cases are $L = 0$

$$\hat{B}_{0,0} = \frac{\hat{N}}{\sqrt{[l]}} = \sum_{m,\nu} a_{m,\nu}^+ a_{m,\nu}, \quad (10)$$

and $L = 1$:

$$\begin{aligned} \hat{B}_{1,0} &= \frac{\sqrt{3}}{\sqrt{l(l+1)[l]}} \hat{L}_z \\ \hat{B}_{1,\pm 1} &= \mp \frac{\sqrt{3}}{\sqrt{2l(l+1)[l]}} \hat{L}_{\pm} \end{aligned} \quad (11)$$

The operator $\hat{B}_{0,0}$ commutes with all the others and the operators $\hat{B}_{L,M}$ for $L \geq 1$ form a Lie-group, which is isomorph to $SU(2l+1)$ with the commutators

$$\begin{aligned} [\hat{B}_{L,M}, \hat{B}_{L',M'}] &= - \sum_{L'',M''} \sqrt{[L][L'] [L'']} \\ &\times [1 - (-1)^{L+L'+L''}] (-1)^{M''} \begin{pmatrix} L & L' & L'' \\ M & M' & -M'' \end{pmatrix} \\ &\times \left\{ \begin{matrix} L & L' & L'' \\ l & l & l \end{matrix} \right\} \hat{B}_{L'',M''} \end{aligned} \quad (12)$$

($\{\dots\}$ denotes the Wigner-6j symbol). Due to the special form of the structure coefficients the operators $\hat{B}_{L,M}$ for odd L form a subgroup, which is isomorph to $SO(2l+1)$. This latter also has a subgroup $SO(3)$ [15] spanned by $\hat{B}_{1,M}$, $M = 0, \pm 1$. The Casimir-operator of $SU(2l+1)$ is

$$\hat{C}_u = \sum_{L=1}^{2l} (-1)^L \hat{B}_L^2 \quad (13)$$

and that of $SO(2l+1)$

$$\hat{C}_o = \sum_{\substack{L=1 \\ L:\text{odd}}}^{2l-1} \hat{B}_L^2. \quad (14)$$

The hamiltonian $\hat{h}$ defined in Eq. (7) commutes with $\hat{L}^2$, $\hat{L}_z$, $\hat{S}^2$, $\hat{S}_z$ and with the Casimir operator $\hat{C}_u$. In fact, the Wigner-3j symbol vanishes for L odd, thus in Eq. (7) the sum over L runs over even values of L . In the special case $l = 1$, because the operator $\hat{B}_2^2 = \hat{C}_u - \hat{C}_o$, and in case $l = 2$, because of the accidental coincidence

$$\begin{pmatrix} 2 & 2 & 2 \\ 0 & 0 & 0 \end{pmatrix}^2 = \begin{pmatrix} 2 & 2 & 4 \\ 0 & 0 & 0 \end{pmatrix}^2 = \frac{2}{35},$$

$\hat{h}$ can be expressed entirely in terms of N , $\hat{C}_u$ and $\hat{C}_o$. For $l > 2$ this is not true anymore, and furthermore $\hat{C}_o$ does not commute with $\hat{h}$.

However, let us consider the model-Hamiltonian

$$\hat{h}_m = \frac{[l]}{8} \hat{N} - \frac{\hat{N}^2}{8} - \frac{[l]}{4(2l+3)} (\hat{C}_u - \hat{C}_o), \quad (15)$$

This model hamiltonian has the following important properties: In the special cases $l = 1, 2$ the two Hamiltonians $\hat{h}$ and $\hat{h}_m$ agree. For $l > 2$ the two hamiltonians are different and we write for the difference

$$\hat{R} \equiv \hat{h} - \hat{h}_m = \sum_{\substack{L=2 \\ L:\text{even}}}^{2l} \left(\begin{pmatrix} l & l & L \\ 0 & 0 & 0 \end{pmatrix}^2 - \frac{2}{(2l+3)[l]} \right) \hat{B}_L^2. \quad (16)$$

In the next sections we show that the averaged value of $\hat{R}$ with the ground states of $\hat{h}_m$ is zero if the particle number equal to

$$N_n = n(2s+1), \quad n = 0, 1, \dots, (2l+1). \quad (17)$$

i.e., N is a multiple of $(2s+1)$. In [10] we have reported that a clusterization in the wave function occur at particle numbers N_n and that the ground state energy per particle has local minima at the n -cluster particle number N_n . We also show that $\hat{h}$ and $\hat{h}_m$ share the same ground states for even particle numbers if $s = 1/2$.

III. CLUSTER STATES

Making a full numerical diagonalization of the hamiltonian (7) is not easy. Fock-vectors have $(2l+1)(2s+1)$ slots, and in each slot there is a zero or 1 due to the fermionic character of the problem. The dimension of the full Hilbert-space is $2^{(2l+1)(2s+1)}$. The spectra do not depend on L_z and S_z , thus we can restrict ourselves to the fermionic sectors $L_z = 0$, $S_z = 0$ (even particle numbers) or $L_z = 0$, $S_z = 1/2$ (odd particle numbers). Conserved operators such as $\hat{S}^2$, $\hat{L}^2$, $\hat{N}$ and $\hat{C}_u$ makes

the numerical problem block-diagonal, but still the computer time and storage required grows exponentially fast as soon as we increase the open shell quantum numbers s or l . This motivates our analytical approach besides the numerical efforts.

In [10] we have investigated the ground state wave function at the particle number N_1 in first quantization. In second quantization the corresponding wave function reads as

$$|1cl\rangle = \sum_k c_k S_m \left(\prod_{i=1}^{(2s+1)} a_{m_i, s+1-i}^+ \right) |0\rangle, \quad (18)$$

$$k \equiv (m_1, \dots, m_{2s+1}),$$

with some coefficients c_k . Here the symbol S_m is an operator which symmetrizes its argument with respect to the indices $m_1, \dots, m_{2s+1}$ (see Sec. IV).

For notational simplicity let us introduce the integer σ by

$$\sigma = \frac{(2s+1)}{2}. \quad (19)$$

The ground state of h_m , explicitly given by

$$|1cl\rangle_0 = S_m(\hat{Q}_{0,0}^{+\sigma})|0\rangle. \quad (20)$$

(see Sec. V). The operator $\hat{Q}_{0,0}^+$ in (20) shall play a central role in the following analysis. It creates a pair state from the vacuum with quantum numbers $L = S = 0$ with fixed open shell quantum numbers (l, s) :

$$\hat{Q}_{0,0} = \frac{1}{\sqrt{[l][s]}} \sum_{m,\nu} (-1)^{s-\nu+l-m} a_{m,\nu}^+ a_{-m,-\nu}^+. \quad (21)$$

Special cases of (18) are

$$|1cl\rangle = |1cl\rangle_0, \quad \begin{cases} l = 1, 2, & s: \text{arbitrary} \\ s = 1/2, & l: \text{arbitrary} \end{cases} \quad (22)$$

since $\hat{h}$ and $\hat{h}_m$ coincides for $l = 1, 2$. The equality for $s = 1/2$ will be considered in Appendix C.

The first example when (22) is not valid occur for $l = 3, s = 3/2$. $N = N_1 = 4$ is the first-cluster particle number. The ground state is in the $L = L_z = S = S_z = 0$ fermionic sector. In this sector there are five orthonormal basis vectors. The matrix elements of $\hat{h}$ on this basis are the following

$$h_{i,j} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{14}{11} & 0 & 0 & 0 \\ 0 & 0 & -\frac{7}{2} & 0 & 0 \\ 0 & 0 & 0 & -\frac{4886}{2145} & -\frac{14}{3} \sqrt{\frac{2}{715}} \\ 0 & 0 & 0 & -\frac{14}{3} \sqrt{\frac{2}{715}} & -\frac{35}{6} \end{pmatrix}$$

The true ground state energy for $N = 4$ comes from the lowest 2×2 block-diagonal and has the value

$$E = -\frac{7(1657 + \sqrt{537729})}{2860} \approx -5.85038,$$

	s=1/2	s=3/2	s=5/2	s=7/2
l=1	-0.75	-3.30	-7.65	-13.80
l=2	-1.25	-4.64	-10.18	-17.86
l=3	-1.75	-5.85	-12.32	-21.17
l=4	-2.25	-6.99	-14.26	-24.12
l=5	-2.75	-8.09	-16.09	—
l=6	-3.25	-9.17	-17.83	—
l=1	-0.75	-3.30	-7.65	-13.80
l=2	-1.25	-4.64	-10.18	-17.86
l=3	-1.75	-5.83	-12.25	-21.00
l=4	-2.25	-6.95	-14.11	-23.72
l=5	-2.75	-8.04	-15.86	—
l=6	-3.25	-9.10	-17.55	—

TABLE I: Ground state energies of $\hat{h}$ (upper part) and $\hat{h}_m$ (lower part) up to two decimal digits precision at the one-cluster particle numbers $N_1 = (2s+1)$.

which is quite close to $h_{5,5} = -35/6 \approx -5.83333$. Numerically $h_{i,j}$ in the same 2×2 block is

$$h_{i,j} \approx \begin{pmatrix} -2.27786 & -0.246813 \\ -0.246813 & -5.83333 \end{pmatrix},$$

and the lowest energy state belongs to the eigenvector

$$v_g \approx \begin{pmatrix} 0.0690865 \\ 1 \end{pmatrix}. \quad (23)$$

It is interesting to present the matrix elements of $\hat{C}_u$

$$C(SU(7))_{i,j} = \begin{pmatrix} \frac{264}{7} & 0 \\ 0 & \frac{264}{7} \end{pmatrix}$$

and that of $\hat{C}_o$

$$C(SO(7))_{i,j} = \begin{pmatrix} 18 & 0 \\ 0 & 0 \end{pmatrix}$$

There is a small admixture of two vectors belonging to different eigenvalues of $\hat{C}_o$ and the dominant vector-part of the eigenvector belong to such a vector, which is an eigenvector of $\hat{C}_o$ with eigenvalue zero. Analyzing further this fifth vector it turns out that it is still given by (20) with $l = 3, s = 3/2$. The average value of $\hat{h}$ by this vector is equal to that of $\hat{h}_m$. Similar property will be proven in Sec. VI also for multi-cluster states. It is important to stress that the fourth vector and the true ground state v_g (23) are also a symmetrized states with $S = 0$.

In Table I. we show the numerically calculated ground state energies of $\hat{h}$ (7) and $\hat{h}_m$ (15). From the data it is clearly seen that for $l = 1, 2$ or for $s = 1/2$ Eq. (22) exact and for all the other cases $|1cl\rangle \approx |1cl\rangle_0$ is a rather good approximation. This shows the importance of the model hamiltonian $\hat{h}_m$. The dominating contribution to

the energy (lower part of Table I) can be written in the form

$$\epsilon_1 = -\sigma\epsilon^{(2)} - \frac{\sigma(\sigma-1)}{2}\delta,$$

where $\epsilon^{(2)}$ and δ are independent of the spin. Their explicit expressions will be derived in Sec. V. The first term gives the energy of σ independent pairs while the second term lowers this energy, indicating clearly that the cluster wave function gives lower energy than the wave function of independent pairs. Note that an eigenfunction of $\hat{h}_m$ exists leading to the eigenvalue $-\sigma\epsilon^{(2)}$ if the inequality $\sigma \leq 2l+1$ is fulfilled. For spin one half particles one has only of course, the first term, the energy of a single pair.

Both $|1cl\rangle$ and $|1cl\rangle_0$ involve the operator S_m , therefore, let us study the properties of the symmetrizer S_m .

IV. THE PROPERTIES OF THE SYMMETRIZER S_m

Let us introduce the symmetrizer symbol S_m by the following properties: *i)* S_m is linear for its argument, *ii)* S_m symmetrizes any pure operator-product $a_{m_1,\nu_1}^+ \dots a_{m_p,\nu_p}^+$ of creation operators a_{m_i,ν_i}^+ , $i = 1, \dots, p$ with respect to *all* m_i without changing the order of spin projections $(\nu_1, \dots, \nu_p)$ including the combinatorial normalization. As an example:

$$S_m(a_{0,3/2}^+ a_{1,1/2}^+ a_{0,-1/2}^+) = \frac{1}{3}(a_{0,3/2}^+ a_{1,1/2}^+ a_{0,-1/2}^+ + a_{1,3/2}^+ a_{0,1/2}^+ a_{0,-1/2}^+ + a_{0,3/2}^+ a_{0,1/2}^+ a_{1,-1/2}^+).$$

(After symmetrization, of course, one can change the order of creation operators as we wish using the anticommutativity of creation operators).

Very important special case is $p = 2s+1$. Writing down all the terms in $S_m(a_{m_1,\nu_1}^+ \dots a_{m_{2s+1},\nu_{2s+1}}^+)$ it turns out that it must be antisymmetric in all spin index $(\nu_1, \dots, \nu_{2s+1})$. Therefore, by inspection we have

$$S_m(a_{m_1,\nu_1}^+ \dots a_{m_{2s+1},\nu_{2s+1}}^+) = \epsilon_{\nu_1, \dots, \nu_{2s+1}} \times S_m(a_{m_1,s}^+ \dots a_{m_{2s+1},-s}^+), \quad (24)$$

where $\epsilon_{\nu_1, \dots, \nu_{2s+1}}$ is the antisymmetric tensor with the convention $\epsilon_{s,s-1, \dots, -s} = 1$. Further expansion is possible for the combination

$$S_m(a_{m_1,s}^+ \dots a_{m_{2s+1},-s}^+) = \frac{1}{(2s+1)!} \sum_{\nu_1} \dots \sum_{\nu_{2s+1}} \epsilon_{\nu_1, \dots, \nu_{2s+1}} a_{m_1,\nu_1}^+ \dots a_{m_{2s+1},\nu_{2s+1}}^+ \quad (25)$$

Next, we enumerate some properties of S_m useful in the

following. If

$$\begin{aligned} \hat{B} &= \sum_{m,n,\nu} f_{m,n} a_{m,\nu}^+ a_{n,\nu} \\ \hat{A}_\alpha &= \sum_{m,n,\nu} (-1)^{s-\nu} g_{m,n}^{(\alpha)} a_{m,\nu}^+ a_{n,-\nu}^+, \quad \alpha = 1, \dots, \sigma, \\ g_{m,n}^{(\alpha)} &= g_{n,m}^{(\alpha)} \end{aligned} \quad (26)$$

where $f_{m,n}$ and $g_{m,n}^{(\alpha)}$ are some numbers, then

$$S_m(\hat{A}_1 \dots \hat{A}_\sigma) = \frac{2^\sigma \sigma! (-1)^{\frac{\sigma(\sigma-1)}{2}}}{(2\sigma)!} \sum_{\substack{m_1, \dots, m_{2\sigma} \\ \nu_1, \dots, \nu_{2\sigma}}} \epsilon_{\nu_1, \dots, \nu_{2\sigma}} \times g_{m_1, m_2}^{(1)} \dots g_{m_{2\sigma-1}, m_{2\sigma}}^{(\sigma)} a_{m_1, \nu_1}^+ a_{m_2, \nu_2}^+ \dots a_{m_{2\sigma}, \nu_{2\sigma}}^+ \quad (27)$$

This identity can be proven using Eqs. (24)-(27) and

$$\begin{aligned} \sum_{\nu_1} \dots \sum_{\nu_\sigma} (-1)^{s-\nu_1} \dots (-1)^{s-\nu_\sigma} \epsilon_{\nu_1, -\nu_1, \dots, \nu_\sigma, -\nu_\sigma} \\ = 2^\sigma \sigma! (-1)^{\frac{\sigma(\sigma-1)}{2}}. \end{aligned} \quad (28)$$

If conditions (26) hold then again from Eqs. (24)-(27) one can derive the identity

$$[\hat{B}, S_m(\hat{A}_1 \dots \hat{A}_\sigma)] = S_m([\hat{B}, \hat{A}_1 \dots \hat{A}_\sigma]), \quad (29)$$

where $[\dots]$ denote a commutator.

V. CLUSTER STATES AND GROUND STATE ENERGIES OF $\hat{h}_m$

Let us consider the states

$$|ncl\rangle_0 = \hat{Q}^{+n}|0\rangle, \quad n = 0, \dots, (2l+1), \quad (30)$$

where the operator $\hat{Q}$ is given by

$$\hat{Q}^+ = S_m(\hat{Q}_{0,0}^{+\sigma}). \quad (31)$$

$\hat{Q}^+$ creates a $(2s+1)$ particle state. In the following we shall prove that $|ncl\rangle_0$ is an eigenstate of the model hamiltonian $\hat{h}_m$ (15). It is easy to show that the state $|ncl\rangle_0$ is annihilated by the Casimir operator $\hat{C}_o$ (See Eq. (14)) of $SO(2l+1)$

$$\hat{C}_o|ncl\rangle_0 = 0 \quad (32)$$

by moving the operators $\hat{B}_{L,M}$ towards to the vacuum using the identity

$$[\hat{B}_{L,M}, S_m(\hat{Q}_{0,0}^{+\sigma})] = \frac{2\sigma(-1)^{L-M}}{\sqrt{[l]}} S_m(\hat{Q}_{L,M}^+ \hat{Q}_{0,0}^{+\sigma-1}), \quad (33)$$

where the operators

$$\hat{Q}_{L,M}^+ = \sqrt{\frac{[L]}{[s]}} \sum_{m=-l}^l \sum_{\nu=-s}^s (-1)^{s-\nu+M} \times \begin{pmatrix} l & l & L \\ m & M-m & -M \end{pmatrix} a_{m,\nu}^+ a_{M-m,-\nu}^+ \quad (34)$$

create also pair states with $S = 0$, but with angular momentum quantum numbers L and M . For $L = M = 0$ this expression agrees with (21). It is easy to prove that

$$\hat{Q}_{2p+1,M} = 0, \quad p : \text{integer}.$$

$\hat{C}_o$ is build up from $\hat{B}_{L,M}$ with L odd but the right hand side of (33) in that case zero, because $\hat{Q}_{L,M}^+ = 0$ for L odd. As a result $\hat{C}_o$ can be moved towards to the vacuum, which is annihilated by $\hat{C}_o$. As a side result we have the property

$$[\hat{C}_o, \hat{Q}^+] = 0, \quad (35)$$

i.e., the operator Q^+ commutes with $\hat{C}_o$.

In order to show that $|ncl\rangle_0$ is also an eigenvector of $\hat{C}_u$ it requires more elaborate calculations. In moving the operators $\hat{B}_{L,M}$ towards the vacuum one encounters several new objects from the commutators. For them one can use the identities (See Appendix A)

$$\sum_{L=0}^{2l} \sum_{M=-L}^L (-1)^M S_m(\hat{Q}_{L,M}^+ \hat{Q}_{L,-M}^+ \hat{Q}_{0,0}^{+\sigma-2}) = [l] S_m(\hat{Q}_{0,0}^{+\sigma}), \quad (36)$$

and

$$\begin{aligned} \sum_{L=0}^{2l} \sum_{M=-L}^L (-1)^M S_m(\hat{Q}_{L,M}^+ \hat{Q}_{0,0}^{+\sigma-1}) S_m(\hat{Q}_{L,-M}^+ \hat{Q}_{0,0}^{+\sigma-1}) \\ = -\frac{[l]}{[s]} S_m(\hat{Q}_{0,0}^{+\sigma})^2. \end{aligned} \quad (37)$$

As a result one has

$$\hat{C}_u |ncl\rangle_0 = \frac{n([l] - n)}{[l]} [s]([l] + [s]) |ncl\rangle_0, \quad (38)$$

i.e., $|ncl\rangle_0$ as given by (30) is really an eigenvector of $\hat{C}_u$. Correspondingly for $\hat{h}_m$ (15) using (32) one obtains:

$$\hat{h}_m |ncl\rangle_0 = \epsilon_n |ncl\rangle_0, \quad (39)$$

with

$$\epsilon_n = -\frac{n(2l+1)(2s+1)}{8(2l+3)} [n(2s-1) + 2l+1 + 4s]. \quad (40)$$

This expression can be cast into the form

$$\epsilon_n = n\epsilon_1 + \frac{n(n-1)}{2} \gamma, \quad (41)$$

where

$$\epsilon_1 = -\frac{(2l+1)(2s+1)}{4(2l+3)} (3s+l) \quad (42)$$

and

$$\gamma = -\frac{(2l+1)(2s+1)}{4(2l+3)} (2s-1). \quad (43)$$

The first term on the right hand side of Eq. (41) can be interpreted as the energy of n independent one-clusters and the second term as a kind of cluster-cluster interaction energy. Furthermore the energy ϵ_1 can be rewritten as

$$\epsilon_1 = -\sigma\epsilon^{(2)} - \frac{\sigma(\sigma-1)}{2} \delta, \quad (44)$$

where

$$\epsilon^{(2)} = \frac{2l+1}{4}, \quad \delta = \frac{12}{2l+3} \epsilon^{(2)} \quad (45)$$

are independent of the spin. For spin one-half particles $\gamma = 0$ and the prefactor of δ in (44) is zero, which means that the clusters consist of pairs which are independent.

One can raise the question what is the ratio of the interaction energies of two pairs within the same cluster and when they belong to two different clusters. This ratio is equal to

$$\frac{\delta}{\gamma/\sigma^2} = \frac{3(2s+1)}{2s-1}, \quad s > 1. \quad (46)$$

It is remarkable that this expression is l -independent. This ratio is always bigger than one, monotonically decreasing with increasing s .

VI. AVERAGE ENERGIES IN CLUSTER STATES

Acting with the operators $\hat{B}_L^2$ (See Eqs. (7),(8)) on the cluster states of $\hat{h}_m$ using the rules (B1),(29) one gets three terms:

$$\begin{aligned} \hat{B}_L^2 |ncl\rangle_0 &= \frac{4n\sigma[l]}{[l]} |ncl\rangle_0 + \frac{4n\sigma(\sigma-1)}{[l]} |\alpha_L\rangle \\ &+ \frac{4n(n-1)\sigma^2}{[l]} |\beta_L\rangle, \end{aligned} \quad (47)$$

where the vectors $|\alpha_L\rangle$ and $|\beta_L\rangle$ are defined as follows:

$$|\alpha_L\rangle = Q^{+(n-1)} \sum_{M=-L}^L (-1)^M S_m(\hat{Q}_{L,M}^+ \hat{Q}_{L,-M}^+ \hat{Q}_{0,0}^{+\sigma-2}) |0\rangle \quad (48)$$

and

$$\begin{aligned} |\beta_L\rangle &= Q^{+(n-2)} \sum_{M=-L}^L (-1)^M \\ &\times S_m(\hat{Q}_{L,M}^+ \hat{Q}_{0,0}^{+\sigma-1}) S_m(\hat{Q}_{L,-M}^+ \hat{Q}_{0,0}^{+\sigma-1}) |0\rangle. \end{aligned} \quad (49)$$

In the special case $L = 0$ the two vectors agree with $|ncl\rangle_0$, and for $L = 2, 4, \dots, 2l$ it can be shown that $|\alpha_L\rangle$ and $|\beta_L\rangle$ can be decomposed as

$$|\alpha_L\rangle = \frac{4[L]}{2l+3}|ncl\rangle_0 + |\alpha_L^\perp\rangle, \quad L = 2, 4, \dots, 2l \quad (50)$$

and

$$|\beta_L\rangle = -\frac{[L]([l] + [s])}{l[s](2l+3)}|ncl\rangle_0 + |\beta_L^\perp\rangle, \quad L = 2, 4, \dots, 2l \quad (51)$$

(See Appendix B), where $|\alpha_L^\perp\rangle$ and $|\beta_L^\perp\rangle$ are possibly zero vectors, but if not, they are eigenvectors of $\hat{C}_o$ with positive eigenvalues, and are automatically orthogonal to $|ncl\rangle_0$ (which is also an eigenvector of $\hat{C}_o$ but with zero eigenvalue).

Using the above results the energy of $\hat{h}$ averaged with $|ncl\rangle_0$ is

$$\frac{\langle ncl|\hat{h}|ncl\rangle_0}{\langle ncl|ncl\rangle_0} = \frac{\langle ncl|\hat{h}_m|ncl\rangle_0}{\langle ncl|ncl\rangle_0} = \epsilon_n, \quad (52)$$

where ϵ_n is given by (40) according to (39).

In Appendix C in the special case $s = 1/2$ we prove the stronger statement (C7), namely that the ground state of $\hat{h}$ is that of $\hat{h}_m$ for even particle numbers. This provides an alternative proof within our framework of Racah's result [11], namely, that the ground state of spin 1/2 particles interacting by an attractive δ -function potential consists of independent pairs.

In order to present some numerical spectra in the most suitable form let us introduce the energies E' defined as

$$E' = E + \frac{Ns(2l+1)}{4}. \quad (53)$$

(53) makes such a shift, proportional to the particle number N , that $E'(N=0) = E'(N=(2l+1)(2s+1)) = 0$, i.e., the energies E' are zero at zero and complete filling. At the cluster numbers N_n the particle number is $N = n(2s+1)$. Making the shift as in (53) the ground state energies E' of the approximate Hamiltonian $\hat{h}_m$ at the cluster number N_n are on the curve

$$E'(N) = \frac{N(2l+1)(2s-1)}{8(2l+3)(2s+1)} [(2s+1)(2l+1) - N]. \quad (54)$$

We show the numerically calculated ground and all excited state energies E' as a function of the particle number N , for $l = 3$ and $s = 1/2$ in Fig. 1, and for $l = 2$ and $s = 3/2$ in Fig. 2.

The spectra are symmetric with respect to the point $N = (2l+1)(2s+1)/2$. This symmetry is a consequence of a particle-hole transformation [10]. Our data show that the lowest energy states at particle numbers N_n are non-degenerate and there is a gap to the first excited states at the same particle numbers.

The regularity in the spectrum in Fig. 2 as compared to the spectrum of Fig. 1 presumably arises from the




FIG. 1: The energy levels E' of the dimensionless hamilton operator as a function of the particle number N for $l = 3$, $s = 1/2$. Individual energy levels are denoted by crosses (+). The dashed line is the function (54).




FIG. 2: The energy levels E' of the dimensionless hamilton operator as a function of the particle number N for $l = 2$, $s = 3/2$. Individual energy levels are denoted by crosses (+). The dashed line is the function (54).

fact that Fig. 2 refers to a situation when the hamilton operator coincides to the model one (15) while this is not the case in Fig. 1.

VII. SUMMARY AND CONCLUSIONS

In preceding sections we have analyzed in great details the clusterization phenomenon in the one open shell model with attractive δ interactions. We have constructed the states $|ncl\rangle_0$ and showed that in a certain subset of shell parameters l, s these states are exact, and in other cases they are quite good approximate ground states. We also showed that there exist a cluster creation operator $\hat{Q}^+$ by which with repeated applications to the vacuum we can create the (approximate or exact) ground

states. We expressed the cluster creation operator $\hat{Q}^+$ in terms of the symmetrized product of the pair creation operator $\hat{Q}_{0,0}^+$ by introducing the symbol S_m .

It is interesting to note that the symbol S_m itself, which by definition symmetrizes with respect to the angular momentum indices, can be interpreted as an antisymmetrizer which act on spin indices. This follows from the form (A2) of the cluster creation operator $\hat{Q}^+ = S_m(\hat{Q}_{0,0}^{+\sigma})$ given by Eq. (31).

Next, let us discuss the main analytic results (52). This equation shows that in the cases $\hat{h} \neq \hat{h}_m$, i.e., when the model and the one open shell Hamiltonian differs, $|ncl\rangle_0$ can be regarded as a variational ansatz, i.e., the ground state energy of the n -cluster state lies below ϵ_n given by (40). In practice $|ncl\rangle_0$ is expected to provide quite a good approximate ground state of $\hat{h}$ as shown in Section III. for a number of examples. Furthermore, if we make the decomposition $\hat{h} = \hat{h}_m + \hat{R}$ and if we consider $\hat{h}_m$ as an unperturbed Hamiltonian with unperturbed ground state $|ncl\rangle_0$ then Eq. (52) shows that we do not obtain correction to the unperturbed energy ϵ_n in first order, corrections are at least of second order, which can be small because the off-diagonal matrix elements can be small. This explains the numerical finding for $l = 3$, $s = 3/2$ and $N = 4$ that the deviation of ϵ_n from the true ground state energy is of the order 0.3% (see Sec. III). The true ground state, in general, is not the state which is annihilated by the Casimir operator of $SO(2l+1)$, but there is a small admixture which mixes some other eigenstates of $\hat{C}_o$ with positive eigenvalues. Note that it remains antisymmetric in the spin indices representing a cluster state.

The main technical problem in deriving values for the approximate (or exact) cluster energies ϵ_n was relegated to Appendix B. Here we applied a powerful projection technics in a non-orthonormal and linearly non-independent basis. This procedure showed that the unperturbed problem for $\hat{h}_m$ is non-degenerate, if we follow how the Casimir of $SO(2l+1)$ acts in the relevant vector space.

APPENDIX A: PROOF OF THE IDENTITIES (36) AND (37)

Let us introduce the spin dependent antisymmetric matrix $\hat{G}_{\nu,z}$ by

$$\hat{G}_{\nu,z} = \sum_m (-1)^{l-m} a_{m,\nu}^+ a_{-m,z}^+. \quad (\text{A1})$$

If we use (27) with $\hat{A}_1 = \dots = \hat{A}_\sigma = \hat{Q}_{0,0}^+$ we have

$$S_m(\hat{Q}_{0,0}^{+\sigma}) = \frac{2^\sigma \sigma! (-1)^{\frac{\sigma(\sigma-1)}{2}}}{(2\sigma)! ([l][s])^{\frac{\sigma}{2}}} \sum_{\nu_1, \dots, \nu_{2\sigma}} \epsilon_{\nu_1, \dots, \nu_{2\sigma}} \times \hat{G}_{\nu_1, \nu_2} \hat{G}_{\nu_3, \nu_4} \dots \hat{G}_{\nu_{2\sigma-1}, \nu_{2\sigma}} \quad (\text{A2})$$

If, on the left hand side of Eq. (36) we use once again (27) with $\hat{A}_1 = \hat{Q}_{L,M}^+$, $\hat{A}_2 = \hat{Q}_{L,-M}^+$ and with $\hat{A}_3 = \dots = \hat{A}_\sigma = \hat{Q}_{0,0}^+$ and perform the standard sum over L and M we arrive to

$$\begin{aligned} & \sum_{L=0}^{2l} \sum_{M=-L}^L (-1)^M S_m(\hat{Q}_{L,M}^+ \hat{Q}_{L,-M}^+ \hat{Q}_{0,0}^{+\sigma-2}) \\ &= \frac{2^\sigma \sigma! (-1)^{\frac{\sigma(\sigma-1)}{2}} [l]}{(2\sigma)! ([l][s])^{\frac{\sigma}{2}}} \sum_{m_1, n_1} (-1)^{l-m_1} (-1)^{l-n_1} \\ &\times \sum_{\nu_1, \dots, \nu_{2\sigma}} \epsilon_{\nu_1, \dots, \nu_{2\sigma}} a_{m_1, \nu_1}^+ a_{n_1, \nu_2}^+ a_{-m_1, \nu_3}^+ a_{-n_1, \nu_4}^+ \\ &\times \hat{G}_{\nu_3, \nu_4} \hat{G}_{\nu_5, \nu_6} \dots \hat{G}_{\nu_{2\sigma-1}, \nu_{2\sigma}} \end{aligned} \quad (\text{A3})$$

If now we change the order of $a_{n_1, \nu_2}^+ a_{-m_1, \nu_3}^+$ using anti-commutativity we can get two more $\hat{G}$ factors. In the next step if we change the second and third indices of the ϵ tensor after comparison with (A2) we arrive to the operator identity Eq. (36).

In order to proof Eq. (37) we proceed as above. Changing the order of two creation operator (and dividing both sides with a common factor) it is left to proof that

$$\begin{aligned} & \frac{1}{[s]} \sum_{\substack{\nu_1, \dots, \nu_{2\sigma} \\ z_1, \dots, z_{2\sigma}}} \epsilon_{\nu_1, \dots, \nu_{2\sigma}} \hat{G}_{\nu_1, \nu_2} \hat{G}_{\nu_3, \nu_4} \dots \hat{G}_{\nu_{2\sigma-1}, \nu_{2\sigma}} \\ & \times \epsilon_{z_1, \dots, z_{2\sigma}} \hat{G}_{z_1, z_2} \hat{G}_{z_3, z_4} \dots \hat{G}_{z_{2\sigma-1}, z_{2\sigma}} \\ &= \sum_{\substack{\nu_1, \dots, \nu_{2\sigma} \\ z_1, \dots, z_{2\sigma}}} \epsilon_{\nu_1, \dots, \nu_{2\sigma}} \hat{G}_{\nu_1, z_1} \hat{G}_{\nu_3, \nu_4} \dots \hat{G}_{\nu_{2\sigma-1}, \nu_{2\sigma}} \\ & \times \epsilon_{z_1, \dots, z_{2\sigma}} \hat{G}_{\nu_2, z_2} \hat{G}_{z_3, z_4} \dots \hat{G}_{z_{2\sigma-1}, z_{2\sigma}}. \end{aligned} \quad (\text{A4})$$

By introducing the spin dependent matrix operator

$$\hat{F}_{\nu,z} = \sum_{\nu_2, \dots, \nu_{2\sigma}} \epsilon_{\nu, \nu_2, \dots, \nu_{2\sigma}} \hat{G}_{z, \nu_2} \hat{G}_{\nu_3, \nu_4} \dots \hat{G}_{\nu_{2\sigma-1}, \nu_{2\sigma}} \quad (\text{A5})$$

and the spin-scalar operator

$$\hat{F} = \sum_{\nu_1, \dots, \nu_{2\sigma}} \epsilon_{\nu_1, \dots, \nu_{2\sigma}} \hat{G}_{\nu_1, \nu_2} \hat{G}_{\nu_3, \nu_4} \dots \hat{G}_{\nu_{2\sigma-1}, \nu_{2\sigma}} \quad (\text{A6})$$

from (A4) it is left to be proven that

$$\frac{1}{2s+1} \hat{F}^2 = \sum_{\nu, z=-s}^s \hat{F}_{\nu,z} \hat{F}_{z,\nu} \quad (\text{A7})$$

Next, we show that

$$\hat{F}_{\nu,z} = \frac{\delta_{\nu,z}}{2s+1} \hat{F}. \quad (\text{A8})$$

If this is true then Eqs. (A7), (A4) and correspondingly (37) are also true.

Let us study first the $\nu \neq z$ case in Eq. (A5). In that case among the summing indices $(\nu_2, \dots, \nu_{2\sigma})$ of the antisymmetric tensor ϵ the index z should occur. Let it

be the index $\nu_i = z$. This cannot be ν_2 , because $\hat{G}_{z,z} = 0$. It means that two *different* $\hat{G}$ has the same index z . However, in the indices ν_2 and ν_{i+1} (if the index z occur in the first index of the second $\hat{G}_{z,\nu_{i-1}}$ or ν_{i-1} (if the index z occur in the second index of the second $\hat{G}_{\nu_{i-1},z}$ factor) the product of two G factor is symmetric, while the antisymmetric tensor ϵ in the same indices is antisymmetric. Summing over ν_2 and ν_{i+1} or ν_{i-1} we get zero. It means, that

$$\hat{F}_{\nu,z} = 0, \quad \text{if } \nu \neq z, \quad (\text{A9})$$

Next, let us study $\hat{F}_{\nu,\nu}$ with ν fixed. This particular index ν should occur among the summing indices of $\hat{F}$. It can be: $\nu_1 = \nu$ or $\nu_2 = \nu$, $\dots$, or $\nu_{2\sigma} = \nu$. Changing the order of indices in all cases such that ν be the first in the ϵ tensor and at the same time in the first $\hat{G}$ factor (Using antisymmetry of the $\hat{G}$ -s and ϵ) all the terms give

$$\begin{aligned} \hat{F} &= (2s+1) \sum_{\nu_2, \dots, \nu_{2\sigma}} \epsilon_{\nu, \nu_2, \dots, \nu_{2\sigma}} \hat{G}_{\nu, \nu_2} \hat{G}_{\nu_3, \nu_4} \dots \hat{G}_{\nu_{2\sigma-1}, \nu_{2\sigma}} \\ &= (2s+1) \hat{F}_{\nu, \nu} \end{aligned} \quad (\text{A10})$$

Eqs. (A9) and (A10) together gives the operator identity (A8), which completes the proof of Eq. (37).

APPENDIX B: DECOMPOSITION OF $|\alpha_L\rangle$ AND $|\beta_L\rangle$

Let us consider how $|\alpha_L\rangle$ or $|\beta_L\rangle$ behaves on applying $\hat{C}_o$ to these vectors. Here we treat the calculation for $|\beta_L\rangle$. Exactly the same method can be applied for $|\alpha_L\rangle$ with one minor difference, which will be shown below.

It is easy to show that

$$\begin{aligned} [\hat{B}_{L_1, M_1}, \hat{Q}_{L_2, M_2}^+] &= 2\sqrt{[L_1][L_2]}(-1)^{M_1} \sum_{\substack{L=0 \\ L:\text{even}}}^{2l} \sum_{M=-L}^M \\ &\times \sqrt{[L]}(-1)^M \begin{pmatrix} L_1 & L_2 & L \\ M_1 & M_2 & -M \end{pmatrix} \begin{Bmatrix} L_1 & L_2 & L \\ l & l & l \end{Bmatrix} Q_{L,M}^+ \end{aligned} \quad (\text{B1})$$

with L_2 even. In using Eqs. (29) and (B1) one obtains

$$\begin{aligned} \hat{C}_o |\beta_L\rangle &= 2[l](1 - \delta_{L,0}) |\beta_L\rangle - 4[L] \sum_{\substack{L_1=0 \\ L_1:\text{even}}}^{2l} \\ &\times \left(\frac{1}{[l]} - \begin{Bmatrix} L & l & l \\ L_1 & l & l \end{Bmatrix} \right) |\beta_{L_1}\rangle \end{aligned} \quad (\text{B2})$$

with L even. In other words, the vector space V spanned by the vectors $\{|\beta_L\rangle | L = 0, 2, \dots, 2l\}$ is invariant under $\hat{C}_o$. If we define matrix elements in V for an operator $\hat{O}$, for which V is an invariant vector space by

$$\hat{O} |\beta_L\rangle = \sum_{L'}' O_{L',L} |\beta_{L'}\rangle, \quad (\text{B3})$$

where $\sum_{L'}'$ stands for $\sum_{L'=0, L':\text{even}}^{2l}$, the matrix elements of $\hat{C}_o$ can be read off (B2):

$$\hat{C}_o |\beta_L\rangle \equiv \sum_{L'}' C_{L',L} |\beta_{L'}\rangle \quad (\text{B4})$$

$$\begin{aligned} C_{L',L} &= 2[l](1 - \delta_{L,0})\delta_{L',L} - 4[L] \left(\frac{1}{[l]} - \begin{Bmatrix} L & l & l \\ L' & l & l \end{Bmatrix} \right) \\ &= (1 - \delta_{L,0})(1 - \delta_{L',0}) \\ &\times \left(2[l]\delta_{L',L} - 4[L] \left(\frac{1}{[l]} - \begin{Bmatrix} L & l & l \\ L' & l & l \end{Bmatrix} \right) \right) \end{aligned} \quad (\text{B5})$$

for L, L' even. The second equality follows from

$$\begin{Bmatrix} L & l & l \\ 0 & l & l \end{Bmatrix} = \begin{Bmatrix} 0 & l & l \\ L & l & l \end{Bmatrix} = \frac{1}{[l]} \quad (\text{B6})$$

for L even. Actually the same matrix elements occur in vector space $V' = \{|\alpha_L\rangle | L = 0, 2, \dots, 2l\}$:

$$\hat{C}_o |\alpha_L\rangle \equiv \sum_{L'}' C_{L',L} |\alpha_{L'}\rangle. \quad (\text{B7})$$

Unfortunately, neither the vectors $|\alpha_L\rangle$, nor the vectors $|\beta_L\rangle$ for $L = 0, 2, \dots, 2l$ are linearly independent. This clearly follows from Eqs. (36) and (37) if we apply both sides to the vacuum and multiply from the left by an appropriate power of $\hat{Q}^+$. However, even in the linearly not independent (and correspondingly not orthonormal) basis one can still calculate matrix elements of operator products such as

$$(\hat{O}^{(1)} \hat{O}^{(2)})_{L',L} = \sum_{L''}' O_{L',L''}^{(1)} O_{L'',L}^{(2)} \quad (\text{B8})$$

provided V is an invariant vector space of $\hat{O}^{(1)}$ and $\hat{O}^{(2)}$, and furthermore, the matrix elements for $\hat{O}^{(1)}$ and $\hat{O}^{(2)}$ are fixed (This statement follows from (B3) if $\hat{O} \equiv \hat{O}^{(2)}$ and we act on both sides with $\hat{O}^{(1)}$ from the left). If one calculates matrix elements of the operator $\hat{R}$

$$\hat{R} = \hat{C}_o^3 - (8l+6)\hat{C}_o^2 + 8l(2l+3)\hat{C}_o \quad (\text{B9})$$

by the well-known properties [14] of the Wigner $6j$ -symbols it turns out that

$$\hat{R}_{L',L} = 0. \quad (\text{B10})$$

It also means that the operator $\hat{C}_o$ on V or V' fulfills

$$\hat{C}_o (\hat{C}_o - 4l\hat{I}) (\hat{C}_o - (4l+6)\hat{I}) = 0, \quad (\text{B11})$$

where $\hat{I}$ is the identity operator.

If an operator $\hat{O}$ has the the minimal polynomial

$$0 = (\hat{O} - \lambda_1 \hat{I}) \dots (\hat{O} - \lambda_d \hat{I}) \quad (\text{B12})$$

with finite d then $\hat{O}$ admits the decomposition

$$\hat{O} = \sum_{i=1}^d \lambda_i \hat{P}_i, \quad (\text{B13})$$

where $\hat{P}_i$ is a projector, i.e., $\hat{P}_i \hat{P}_j = \hat{P}_i \delta_{i,j}$ and

$$\hat{P}_i = \prod_{\substack{j=1 \\ j \neq i}}^d \frac{\hat{O} - \lambda_j \hat{I}}{\lambda_i - \lambda_j} \quad (\text{B14})$$

From Eq. (B11) is clear that on V at most we have $d = 3$, and the three eigenvalues of $\hat{C}_o$ are $\lambda_1 = 0$, $\lambda_2 = 4l$ and $\lambda_3 = (4l+6)$ respectively. Our main purpose is to calculate the orthogonal projection of $|\beta_L\rangle$ to $\hat{Q}^{+n}|0\rangle = |\beta_0\rangle$. This latter vector is an eigenvector of $\hat{C}_o$ with eigenvalue 0, thus let us consider the projector of $\hat{P}_1$:

$$\hat{P}_1 = \frac{(\hat{C}_o - 4l\hat{I})(\hat{C}_o - (4l+6)\hat{I})}{4l(4l+6)}. \quad (\text{B15})$$

Taking matrix elements on both sides is easy. Straight-forward calculation leads to

$$\left(\hat{P}_1\right)_{L',L} = \delta_{L,0}\delta_{L',0} + \frac{[L]}{l(2l+3)}(1 - \delta_{L,0})(1 - \delta_{L',0}) \quad (\text{B16})$$

If we consider the decomposition

$$|\beta_L\rangle = \hat{P}_1|\beta_L\rangle + (\hat{I} - \hat{P}_1)|\beta_L\rangle \quad (\text{B17})$$

the first term $\hat{P}_1|\beta_L\rangle$ is vector which is an eigenvector of $\hat{C}_o$ with eigenvalue zero, the second term, if it is nonzero, however belong to the subspace in which $\hat{C}_o$ has positive eigenvalues. Thus, the two vectors on the right hand side of (B17) are orthogonal to each other. Most important is the first term. Knowing the matrix elements of $\hat{P}_1$ it reads as

$$\hat{P}_1|\beta_L\rangle = \delta_{L,0}|\beta_0\rangle + \frac{[L](1 - \delta_{L,0})}{l(2l+3)} \sum_{\substack{L'=2 \\ L':\text{even}}}^{2l} |\beta_{L'}\rangle \quad (\text{B18})$$

Equation (37) implies (by applying both side to the vacuum and multiplying by $\hat{Q}^{+(n-2)}$ from the left)

$$\sum_{\substack{L'=2 \\ L':\text{even}}}^{2l} |\beta_{L'}\rangle = -\frac{[l] + [s]}{[s]} |\beta_0\rangle \quad (\text{B19})$$

Putting Eqs. (B17)-(B19) together we obtain the anticipated result of Eq. (51)

To prove the corresponding result (50) for the decomposition of $|\alpha_L\rangle$ we can proceed as above, but instead of (B19) we should use

$$\sum_{\substack{L'=2 \\ L':\text{even}}}^{2l} |\alpha_{L'}\rangle = 2l|\alpha_0\rangle, \quad (\text{B20})$$

which follows from the operator identity (36) if we apply both sides to the vacuum and multiply by $\hat{Q}^{+(n-1)}$ from the left.

APPENDIX C: THE SPECIAL CASE $s = 1/2$

In the special case $s = 1/2$ the parameter σ is equal to 1 and because of the only two possibilities for $\nu = 1/2 = \uparrow$ or $\nu = -1/2 = \downarrow$ there is no need to symmetrize $\hat{Q}_{L,M}^+$ as defined in (34)

$$S_m(\hat{Q}_{L,M}^+) = \hat{Q}_{L,M}^+. \quad (\text{C1})$$

Consequently, according to (31)

$$\hat{Q}^+ = \hat{Q}_{0,0}^+. \quad (\text{C2})$$

By definition Eq. (34) gives for $s = 1/2$

$$\begin{aligned} \hat{Q}_{L,M}^+ &= \sqrt{2[L]}(-1)^M \\ &\times \sum_{m=-l}^l \begin{pmatrix} l & l & L \\ m & M-m & -M \end{pmatrix} a_{m,\uparrow}^+ a_{M-m,\downarrow}^+. \end{aligned} \quad (\text{C3})$$

Acting with $\hat{h}$ to $|ncl\rangle_0$ Eqs. (7), (47) and (40) for $s = 1/2$ leads to

$$\hat{h}|ncl\rangle_0 = \epsilon_n|ncl\rangle_0 - \frac{[l]}{2}n(n-1) \sum_{L=0}^{2l} \begin{pmatrix} l & l & L \\ 0 & 0 & 0 \end{pmatrix}^2 |\beta_L\rangle. \quad (\text{C4})$$

Next we show that the second term is identically zero for $s = 1/2$. Using (C2) (C3) and Eq. (5) it is easy to check that

$$\begin{aligned} \sum_{L=0}^{2l} \begin{pmatrix} l & l & L \\ 0 & 0 & 0 \end{pmatrix}^2 |\beta_L\rangle &= \frac{8\pi}{[l]^2} \hat{Q}_{0,0}^{+(n-2)} \sum_{\substack{m_1, m_2 \\ m_3, m_4}} (-1)^{m_1+m_2} \\ &\times f_{m_1, m_2; m_3, m_4} a_{m_1, \uparrow}^+ a_{m_2, \downarrow}^+ a_{m_3, \uparrow}^+ a_{m_4, \downarrow}^+ |0\rangle \end{aligned} \quad (\text{C5})$$

If we insert into this equation the second form (6) for f we arrive at the rather long form

$$\begin{aligned} \sum_{L=0}^{2l} \begin{pmatrix} l & l & L \\ 0 & 0 & 0 \end{pmatrix}^2 |\beta_L\rangle &= 2\hat{Q}_{0,0}^{+(n-2)} \sum_{\substack{m_1, m_2 \\ m_3, m_4}} (-1)^{m_1+m_3} \\ &\times \sum_{\substack{L=0 \\ L:\text{even}}}^{2l} [L] \begin{pmatrix} l & l & L \\ 0 & 0 & 0 \end{pmatrix}^2 \sum_{M=-L}^L \begin{pmatrix} l & l & L \\ m_1 & m_3 & -M \end{pmatrix} \\ &\times \begin{pmatrix} l & l & L \\ -m_4 & -m_2 & -M \end{pmatrix} a_{m_1, \uparrow}^+ a_{m_2, \downarrow}^+ a_{m_3, \uparrow}^+ a_{m_4, \downarrow}^+ |0\rangle \end{aligned} \quad (\text{C6})$$

The operator $a_{m_1, \uparrow}^+ a_{m_2, \downarrow}^+ a_{m_3, \uparrow}^+ a_{m_4, \downarrow}^+$ is antisymmetric in m_1, m_3 , while its coefficient is symmetric in the same indices, thus, the right hand side is zero. this completes the proof that the state $|ncl\rangle_0$ is an eigenstate of $\hat{h}$:

$$\hat{h}|ncl\rangle_0 = \epsilon_n|ncl\rangle_0, \quad s = 1/2, \quad (\text{C7})$$

with

$$\epsilon_n = -\frac{n(2l+1)}{4}, \quad s = 1/2. \quad (\text{C8})$$

ACKNOWLEDGMENTS

Nos. OTKA T046129 and T038202.

The present work has been partially supported by the Hungarian Research National Foundation under Grant

-
- [1] U. Fano, G. Racah, *Irreducible Tensorial Sets*, (Academic Press, New York, 1959).
 - [2] B. R. Judd, *Second Quantization and Atomic Spectra*, (The John Hopkins Press, Baltimore, 1967).
 - [3] L. Szasz, *The Electronic Structure of Atoms*, (John Wiley & Sons Inc., New York, 1992).
 - [4] P. Ring and P. Schuck, *The Nuclear Many-Body Problem*, 3. printing (Springer, Berlin, 2004).
 - [5] C. J. Pethick, H. Smith *Bose-Einstein Condensation of Dilute Gases*, (Cambridge University Press, Cambridge, 2004)
 - [6] H. T. C. Stoof and M. Houbiers, Proceedings of the International School of Physics - Enrico Fermi, eds. M. Inguscio, S. Stringari and C.E. Wieman, (IOS Press, 1999).
 - [7] See Reference [5] Chapter 14.
 - [8] Q. Chen, J. Stajic, S. Tan, and K. Levin, Physics Reports **412**, 1 (2005).
 - [9] H. Heiselberg, New J. Physics **6**, 137 (2004).
 - [10] A. Csordás, P. Szépfalusy, É. Szőke, Phys. Rev. Lett. **92**, 090401 (2004).
 - [11] G. Racah, Physica **18**, 1097 (1952).
 - [12] G. M. Bruun and H. Heiselberg, Phys. Rev. A **65**, 053407 (2002).
 - [13] G. M. Bruun, Phys. Rev. A **66**, 041602(R) (2002).
 - [14] A. R. Edmonds, *Angular Momentum in Quantum Mechanics*, (Princeton University Press, Princeton, N. J., 1957).
 - [15] M. Hamermesh, *Group Theory and its Application to Physical Problems*, (Addison-Wesley Publishing Company Inc., Reading, Massachusetts, 1964).